

Topex® Handicaine Topical Anesthetic Gel – 20% Benzocaine

DIRECTIONS FOR USE

REF AD31023 – Handicaine Stix, Cherry, Box of 24
REF AD31024 – Handicaine Stix, Cherry, Box of 120
REF AD31025 – Handicaine Stix, Pina Colada, Box of 24
REF AD31026 – Handicaine Stix, Pina Colada, Box of 120
NPN Number: 00638358

For dental use only.

1. DESCRIPTION

Flavored 20% benzocaine gel for topical mucosal anesthesia, delivered in an easy-to-use single dose package with applicator.

2. INDICATIONS

Indicated as a topical anesthetic for use on oral mucosa prior to local anesthetic injections, scaling and prophylaxis. Also useful to relieve discomfort associated with taking impressions and intra-oral radiographs.

3. CONTRAINDICATIONS

Should not be used with individuals with a known sensitivity to benzocaine or PABA.

4. WARNINGS

Keep out of reach of children. For professional dental use only.

5. PRECAUTIONS

Topex Handicaine Stix are intended for single patient use. Do not attempt to reuse the applicator or refill the tray.

6. DOSAGE AND ADMINISTRATION

1. Each Topex Handicaine Stix dose contains approximately 0.4g of 20% benzocaine anesthetic gel (approximately 7-9mg benzocaine per unit dose tray).
2. Open unit dose tray by peeling foil lid stock from the plastic tray. Dispose of foil.
3. Using the included cotton applicator, apply the benzocaine anesthetic gel to the mucosa to achieve topical anesthesia.
4. For topical tissue anesthesia during scaling procedures, coat the scaler with the product prior to use.

7. STORAGE

Store between 59°-86°F (15°-30°C). Protect from freezing.



Sultan Healthcare
1301 Smile Way • York, PA 17404-0807 • USA
Toll Free: 800-637-8582 • Phone: 201-871-1232
Fax: 201-871-0321 • www.sultanhealthcare.com

Made in the USA
0030501DF, R1-091014