

Topex® Topical Anesthetic Gel – 20% Benzocaine

DIRECTIONS FOR USE

Gel – 1oz/30mL

REF AD31001 Cherry
REF AD31002 Mint
REF AD31004 Pina Colada
REF AD31005 Bubble-Num
REF AD31006 Strawberry
REF AD31007 Razzberry

Gel – 1oz/30mL

REF CN31001FG Cherry
REF CN31002FG Mint
REF CN31004FG Pina Colada
REF CN31005FG Bubble-Num
REF CN31006FG Strawberry
REF CN31007FG Razzberry

Gel – 100g jar

REF AD31064 Cherry
REF AD31065 Mint
REF AD31067 Pina Colada
REF AD31068 Bubble-Num

NPN Number: 00638358

For dental use only.

1. DESCRIPTION

Flavored 20% benzocaine gel for topical mucosal anesthesia.

2. INDICATIONS

Indicated as a topical anesthetic for use on oral mucosa prior to local anesthetic injections, scaling and prophylaxis. Also useful to relieve discomfort associated with taking impressions and intra-oral radiographs.

3. CONTRAINDICATIONS

Should not be used with individuals with a known sensitivity to benzocaine or PABA.

4. WARNINGS

Keep out of reach of children. For professional dental use only.

5. PRECAUTIONS

Topex 20% benzocaine gels is supplied in multiple use containers. Take care not to contaminate the bottle by reintroducing a used cotton applicator into the bottle.

6. DOSAGE AND ADMINISTRATION

1. Each gram of Topex 20% benzocaine gel contains between 180-220mg benzocaine in a flavored base.
2. Using a new cotton applicator, apply a small amount of gel to the mucosa to achieve topical anesthesia. Do not reintroduce the cotton applicator into the bottle.
3. For topical tissue anesthesia during scaling procedures, dispense a small amount of Topex gel into a dappen dish, then coat the scaler with the product prior to use.
4. Tightly re-cap the jar after each use.

7. STORAGE

Store between 59°-86°F (15°-30°C). Protect from freezing.



Sultan Healthcare
1301 Smile Way • York, PA 17404-0807 • USA
Toll Free: 800-637-8582 • Phone: 201-871-1232
Fax: 201-871-0321 • www.sultanhealthcare.com

Made in the USA
0031001DF, R2-102814